

Instructions: Keep all devices capable of communication turned off and out of sight. The exam lasts for 1 hour and 15 min. **Multiple Choice Questions (65 points)**

Q1 Evaluate $\int_0^2 \int_1^x x^2 y \, dy \, dx$

(A) 28/15 (C) 1/24 (E) None

(B) 3 (D) 1/13

Q2 Evaluate $\int_0^e \int_0^e \int_0^e \frac{1}{xyz} \, dx \, dy \, dz$

(A) π (C) e (E) None

(B) $\ln 2 + 1$ (D) $\sqrt{2}e$

Q3 Evaluate the line integral

$$\oint_C (x^2 + y^2)dx + 2xydy,$$

where C is the square bounded by the lines $x = 0, x = 2, y = 0, y = 2$.

(A) -1 (C) $1/2$ (E) None

(B) 0 (D) 13

Q4 Find the directional derivative of the function $g(x, y) = (x + 3y)^2$

at the given point $(1, -1)$ in the direction of the vector $\vec{v} = \frac{1}{\sqrt{2}}(1, -1)$.

(A) $\sqrt{2} + 2$ (C) $4\sqrt{2}$ (E) None

(B) $\sqrt{3} - 1$ (D) 5

Q5 Let $z = 4 + x^3 + y^3 - 3xy$. Which of the following statements are true?

1. $(1, 1)$ is a local maximum,

2. $(0, 0)$ is a saddle point

3. $(2, 4)$ is a local minimum.

(A) Only 1 (C) Only 3 (E) None

(B) Only 2 (D) 1 and 3

Q6 Find $\lim_{t \rightarrow 0} \left[\frac{\sin t}{t} \vec{i} + t \vec{j} + (t-1)^4 \vec{k} \right]$

(A) \vec{i} (C) $\vec{i} + \vec{j} + \vec{k}$ (E) None

(B) $\vec{i} + \vec{j}$ (D) $\vec{i} - \vec{k}$

Q7 What is the length of the arc described $r(t) = (3t^2 \vec{i} + 2t^3 \vec{j} + \vec{k})$

(A) $4\sqrt{2}$ (C) $4\sqrt{2} - 2$ (E) None

(B) $4\sqrt{2} + 1$ (D) $\sqrt{2} + 4$

Q8 Compute $\int_{-1}^1 \sqrt{1-x^2} dx$

(A) $\frac{\pi}{2}$ (C) π^2 (E) None

(B) 2π (D) $2\pi^2$

Q9 Evaluate the series $\sum_{n=1}^{\infty} 2^{4-3n}$

(A) 125/9 (C) 63/7 (E) None

(B) 161/5 (D) 128/7

Q10 Find the radius of convergence of the power series $\sum_{n=1}^{\infty} \frac{n^3 x^n}{3^n}$

(A) 1 (C) 3 (E) None

(B) 1.5 (D) 4

Q11 The value of $\lim_{(x,y,z) \rightarrow (2,3,0)} [xe^z + \ln(2x-y)]$

(A) 2 (C) 3 (E) None

(B) $2\sqrt{2}$ (D) DNE

Q12 Compute $\int \frac{dx}{x^2 - x}$.

(A) $\ln \left| \frac{x-1}{x} \right| + C$ (C) $\frac{1}{2} \ln |x^2 + 1| + C$

(B) $2 \ln |x| + \frac{1}{3} \ln |x^2 - x| + C$ (D) None

Q13 Compute $\int_0^1 x e^x dx$.

(A) -1 (C) 1 (E) None

(B) 0 (D) 2

True and False

Q1 The surface $x^2 + y^2 + z^2 = 2z$ is a sphere.

T

F

Q2 The vector $(1, 2, 3)$ is perpendicular to the plane $2x + 4y + 6z = 4$.

T

F

Q3 $(0, 0)$ is a local minimum of the function $f(x, y) = x^6 + y^6$.

T

F

Q4 If $|\vec{v} \times \vec{w}| = 0$ then $\vec{v} = 0$ or $\vec{w} = 0$.

T

F

Q5

T

F

Q6 If the series $\sum_{n=1}^{\infty} a_n$ converges, then the sequence a_n converges to 0.

T

F

Q7 The $\sum_{n=1}^{\infty} \frac{1}{n^2}$ is convergent.

T

F

Q8 The $\int_1^{\infty} \frac{1}{x^2} dx$ is divergent.

T

F

Q9 The intersection of the sphere $x^2 + y^2 + z^2 = 169$ with the plane $z = 14$ is a circle.

T

F

Q10 The vectors $\vec{v} = (2, 1, 5)$ and $\vec{w} = (2, 1, -1)$ are perpendicular.

T

F

Fill in the blanks

Q1 The _____ from the point $P = (x_1, y_1, z_1)$ to the _____ $ax + by + cz + d = 0$ is given by

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}.$$

Q2 If $\|\vec{v}\| = 1$ then \vec{v} is called _____

Q3 A point (a, b) in the plane is called a _____ of a function $f(x, y)$ if $\nabla f(a, b) = (0, 0)$.

Q4 A _____ of radius R centered at $P = (a, b, c)$ is the collection of points in space which have distance R from P .

Q5 If f_{xy} and f_{yx} are both _____ then $f_{xy} = f_{yx}$.

Classical Problems . Show all your work. No work=No credit!


Q3(10pts) Find an equation for the plane passing through the origin that is perpendicular to the tangent plane to the graph of

$$z = f(x, y) = 3x^2 - y^2 - 3x + 3y$$

at the point $(1, -1, f(1, -1))$.

Solution:

Q2 Compute the line integral $\int_C xy^2 dx - x^2 y dy$ by applying Green's theorem.



Solution: